

Technical Data Sheet



InVivoMAb anti-mouse LPAM-1 (Integrin $\alpha 4\beta 7$)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: **BE0034**
Clone: **DATK32**
Isotype: Rat IgG2a, κ
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: TK1 cells
Reported Applications: *in vivo* Integrin $\alpha 4\beta 7$ neutralization
Flow cytometry
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_1107713](https://eutils.ncbi.nlm.nih.gov/entrez/eutils/rrid.cgi?db=AB)
Molecular Weight: 150 kDa

Description

The DATK32 monoclonal antibody reacts with mouse LPAM-1 also known as integrin alpha 4 beta 7. The 130 kDa integrin $\beta 7$ chain associates with the 150 kDa integrin $\alpha 4$ (CD49d) chain to form LPAM-1, a member of the Ig superfamily. LPAM-1 is expressed by peripheral lymphocytes, small subsets of thymocytes, and bone marrow progenitors. LPAM-1 binds VCAM-1 (CD106), MAdCAM-1, and fibronectin and facilitates lymphocyte adhesion and migration to the intestine and associated lymphoid tissues. The DATK32 antibody has been reported to block LPAM-1-mediated cell adhesion *in vivo*. Bio X Cell is pleased to offer a recombinant, murine chimeric version of the original DATK32 antibody, DATK32-CP069. The variable domain sequences are identical to DATK32, but the constant region sequences have been switched from rat IgG2a, κ to mouse IgG2a, κ for use in murine models. Species-matched chimeric antibodies exhibit regulated effector functions—including Fc receptor binding and complement activation—and result in less immunogenicity and formation of anti-drug antibodies (ADAs) than xenogenic antibodies in animal models. The highly controlled sequence and lack of genetic drift in recombinant antibodies provide more reliable and reproducible results over hybridoma derived antibodies.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0034?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC