

# Technical Data Sheet

## InVivoMAb anti-mouse GM-CSF



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

Lot Number: Lot Specific\*  
Volume: Lot Specific\*  
Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*  
Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

Catalog Number: BE0259  
Clone: MP1-22E9  
Isotype: Rat IgG2a,  $\kappa$   
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol  
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer  
Immunogen: Recombinant mouse GM-CSF  
Reported Applications: *in vivo* GM-CSF neutralization  
*in vitro* GM-CSF neutralization  
Flow cytometry  
Formulation: PBS, pH 7.0  
Contains no stabilizers or preservatives  
Endotoxin: <2EU/mg (<0.002EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
Purity: >95%  
Determined by SDS-PAGE  
Sterility: 0.2  $\mu$ m filtered  
Production: Purified from cell culture supernatant in an animal-free facility  
Purification: Protein G  
RRID: [AB\\_2687738](https://www.ebi.ac.uk/ols/ontologies/ab/term/AB_2687738)  
Molecular Weight: 150 kDa

### Description

The MP1-22E9 monoclonal antibody reacts with mouse granulocyte-macrophage colony-stimulating factor (GM-CSF), also known as colony stimulating factor 2 (CSF2). GM-CSF is a 14 kDa monomeric hematopoietic factor secreted by macrophages, T cells, mast cells, NK cells, endothelial cells and fibroblasts. GM-CSF stimulates stem cells to differentiate into granulocytes (neutrophils, eosinophils, and basophils) and monocytes. The MP1-22E9 antibody is a GM-CSF neutralizing antibody.

### Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/be0259?bxcs=9k1b3a#tab\\_references](https://bioxcell.com/be0259?bxcs=9k1b3a#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC**