Technical Data Sheet

InVivoMAb anti-mouse IL-12 p35

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0371**
Clone:	**C18.2**
Isotype:	Rat IgG2a, κ
Recommended Isotype Control(s):	InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Recombinant mouse IL-12 p70
Reported Applications:	*in vivo* IL-12 p35 neutralization *in vitro* IL-12 p35 neutralization ELISA Immunoprecipitation
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 μm filtration
Purification:	Protein G
RRID:	AB_2927508
Molecular Weight:	150 kDa

Description

The C18.2 monoclonal antibody reacts with mouse interleukin-12 (IL-12) p35. IL-12 is a heterodimeric 70 kDa cytokine consisting of two covalently linked subunits, 40 kDa (p40) and 35 kDa (p35). IL-12 is secreted by activated monocytes, macrophages, and dendritic cells in response to bacterial pathogens or products such as lipopolysaccharides (LPS). IL-12 is a potent regulator of cell-mediated immune responses and plays a key role in the development of Th1 responses, leading to IFNγ and IL-2 production.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0371?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com