

InVivoMAb anti-mouse CD31 (PECAM-1)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0377
Clone: 390
Isotype: Rat IgG2a, κ
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: C3H/HeJ mouse hematopoietic progenitor cell line 3
Reported Applications: *in vivo* CD31 blocking
Intravital imaging
Immunofluorescence
Flow cytometry
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtration
Purification: Protein G
RRID: [AB_2927514](https://abnova.com/AB_2927514)
Molecular Weight: 150 kDa

Description

The 390 monoclonal antibody reacts with mouse CD31, also known as PECAM-1 (platelet endothelial cell adhesion molecule). CD31 is a 130-140 kDa glycoprotein and member of the immunoglobulin superfamily. CD31 is expressed constitutively on the surface of adult and embryonic endothelial cells and is weakly expressed on platelets, granulocytes, monocytes/macrophages, dendritic cells, and T and B cell subsets. CD31 mediates homophilic and heterophilic cell-cell adhesion and is critical for cell-cell interactions. Ligands for CD31 are reported as CD38 and vitronectin receptor (α v β 3 integrin, CD51/CD61). The 390 antibody has been shown to inhibit *in vivo* functions mediated by CD31.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0377?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC