

Technical Data Sheet

InVivoMAb anti-mouse MHC Class I (H-2Db)



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0451
Clone: 28-14-8S
Isotype: Mouse IgG2a, κ
Recommended Isotype Control(s): InVivoMAb mouse IgG2a isotype control, unknown specificity
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: C3H.SW mouse splenocytes
Reported Applications: MHC-I Immunopeptidomics
in vitro blocking of MHC-I
Immunocapture of peptide-MHC class - I complexes
Immunoprecipitation
Flow cytometry
Immunohistochemistry (frozen)
ELISA
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID:
Molecular Weight: 150 kDa

Description

The 28-14-8S monoclonal antibody reacts with the $\alpha 3$ domains of H-2Db of the MHC class I alloantigen expressed on nucleated cells from mice of the H-2Db haplotype. The 28-14-8S antibody shows weak cross-reactivity with Ld antigen, but it doesn't cross-react with Dd. H-2Db is a member of the MHC subgroup, and it is characterized by a hydrophobic ridge in the binding cleft. The H2 class I molecules are made up of three parts: a heavy chain that is 42 kDa, a 12-kDa protein called $\beta 2$ -microglobulin, and a short peptide made up of 8 to 11 amino acids. The endoplasmic reticulum carries out this assembly, but the stable trimeric complexes translocate to cell surfaces. H-2Db is a class I molecule that is able to reach the cell surface even in the absence of either $\beta 2$ microglobulin or TAP-provided peptides. Interestingly, despite being incomplete and unstable, this molecule is still able to exhibit some functional capacities, in that it can induce and be recognized by allogeneic and self-restricted cytotoxic T lymphocytes. The H-2Db plays a critical role in antigen presentation to T cells expressing CD3/TCR and CD8 proteins. Like other MHC antibodies, the 28-14-8S clone finds its application in various immunoassays involving MHC-peptide complexes. It is a useful tool for immunopeptidomics research, making it easier to identify and describe neoantigens using HPLC and mass spectrometry.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0451?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC