

Technical Data Sheet

FlowMAb FITC anti-mouse Ly6G



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: FM0075-1-FITC
Clone: 1A8
Isotype: Rat IgG2a, κ
Conjugation: FITC
Excitation Source: Blue 488 nm
Excitation Max: 494 nm
Emission Max: 518 nm
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: EL4J cells transfected with Ly6G
Reported Applications: *in vivo* neutrophil depletion
in vivo MDSC depletion
Immunofluorescence
Immunohistochemistry (paraffin)
Immunohistochemistry (frozen)
Flow cytometry
Formulation: PBS, pH 7.0
Contains 0.09% Sodium Azide
Endotoxin: <2EU/mg (<0.002EU/μg)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μm filtration
Purification: Protein G
RRID: [AB_1107721](https://abnova.com/AB_1107721)
Molecular Weight: 150 kDa

Description

The 1A8 monoclonal antibody reacts with mouse Ly6G. Ly6G is a 21-25 kDa member of the Ly-6 superfamily of GPI-anchored cell surface proteins with roles in cell signaling and cell adhesion. Ly6G is expressed differentially during development by cells in the myeloid lineage including monocytes, macrophages, granulocytes, and neutrophils. Monocytes typically express Ly6G transiently during development while mature granulocytes and peripheral neutrophils retain expression making Ly6G a good cell surface marker for these populations. Unlike the RB6-8C5 antibody, the 1A8 antibody reacts specifically with mouse Ly6G with no reported cross reactivity with Ly6C.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.


Application References

For a complete list of references, visit https://bioxcell.com/fm0075-1-fitc?bxcs=9k1b3a#tab_references or scan the QR code below.



Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC
<https://bioxcell.com>
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC